arXiv:1505.06904v1 [math.CA] 26 May 2015

Choldowsky type generalization of the $q-$Favard-Szàsz operators

Ali Karaisa *

Department of Mathematcs-Computer Sciences, Faculty of Science, Necmettin Erbakan Unversity Meram
Campus , 42090 Meram, Konya, Turkey

Abstract. In the present paper, we introduce a Choldowsky type generalization of the $q-$Favard-Szàsz operators and obtain weighted statistical approximation properties of these operators. We also establish the rates of statistical convergence by means of the modulus of continuity and the Lipschitz type maximal function. Further, we study the local approximation properties of these operators.

Keywords: Choldowsky type generalization of the $q-$Favard-Szàsz operators, modulus of continuity, local approximation, Peetre's K-functional, statistical convergence.

MSC:41A25, 41A36

1. Introduction

The approximation of function by using the linear positive operators introduced via $q-$ calculus is currently be come active research. After the paper of Phillips [6] who generalized the classical Bernstein polynomials base on $q-$integer, many generalization of well-known positive linear operators, base on $q-$integer were introduce and studied several authors. We refer to the following studies related to the approximation properties of the $q-$analogues of some operators, [1–4]. Recently, the statistical approximation properties have been invesigated for $q-$analogue polynomials. For example in [12, 13] $q-$analogue of Kantorovich-Bernstein operators; in [14] $q-$Baskakov-Kantorovich operators; [15] $q-$Szàsz-Mirakjan operators; in [16] modified $q-$Stancu-Beta operators were introduced and their statistical properties were investigated. For further information related to the statistical approximation of the operators, the followings are remarkable among others [18–21].

In this work, we generalized a Choldowsky type Favard-Szàsz operators base on $q-$integer and we study the weighted statistical approximation properties of the Choldowsky type $q-$Favard-Szàsz operators via Korovkin type approximation theorem. Further we compute the rate of statistical convergence by using modulus of continuity. Furthermore we also obtain some local approximation results of these new operators.

First of all, we recall some definitions and notations regarding the concept of $q-$calculus. For any none-negative integer r, the $q-$ integer of the number r is defied by

$$[r]_q := \begin{cases} \frac{1-q^r}{1-q}, & \text{if } q \neq 1 \\ 1, & \text{if } q = 1 \end{cases}.$$

The $q-$factorial $[n]_q!$ and $q-$binomial coefficients are defined as

$$[n]_q! := \begin{cases} [n]_q [n-1]_q \cdots [1]_q, & n \in \mathbb{N} \\ 1, & n = 0 \end{cases}$$

and

$$\left[\begin{array}{c} n \\ k \end{array} \right]_q = \frac{[n]_q!}{[k]_q! [n-k]_q!}, 0 \leqslant k \leqslant n.$$

The $q-$derivative $D_q f$ of a function f is defined by

$$(D_q f)(x) = \frac{f(x) - f(qx)}{(1-q)x},\ x \neq 0.$$

Also, if there exists $\frac{df}{dx}(0)$, then $(D_q f)(0) = \frac{df}{dx}(0)$. The following $q-$derivatives of the product of the functions $f(x)$ and $g(x)$ are equivalent:

$$D_q(f(x)g(x)) = f(qx) D_q g(x) + g(x) D_q f(x)$$

and

$$D_q(f(x)g(x)) = f(x) D_q g(x) + g(qx) D_q f(x).$$

The $q-$analogues of the exponential function are given by

$$e_q^x = \sum_{n=0}^{\infty} \frac{x^n}{[n]_q!}$$

---

[*] Corresponding author: Tel:+90 332 323 8220; fax:+90 332 323 8245 E-mail:akaraisa@konya.edu.tr

and

$$E_q^x = \sum_{n=0}^{\infty} q^{\frac{n(n-1)}{2}} \frac{x^n}{[n]_q!}.$$

The exponential functions have the following properties:

$$D_q\left(e_q^{ax}\right) = a e_q^{ax},\ D_q\left(E_q^{ax}\right) = a E_q^{aqx},\ e_q^x E_q^{-x} = E_q^x e_q^{-x} = 1.$$

Further results can be found in [8].

2. Construction of the operators

In [9], Jakimovski and Leviatan introduced a Favard Szàsz type operator, by using Appell polynomials $p_k(x) \geqslant 0$ defined by

$$g(u)\, e^{-ux} = \sum_{k=0}^{\infty} p_k(x)\, u^k,$$

where $g(z) = \sum\limits_{n=0}^{\infty} a_n z^n$ is an analytic function in the disc $|z| < R$, $R > 1$ and $g(1) \neq 0$,

$$P_{n,t}(f;x) = \frac{e^{-nx}}{g(1)} \sum_{k=0}^{\infty} p_k(nx)\, f\left(\frac{k}{n}\right)$$

and they investigated some approximation properties of these operators.

Atakut at all. [7] defined a Choldowsky type of Favard-Szàsz operators as follows:

$$P^*(f;x) = \frac{e^{-\frac{nx}{b_n}}}{g(1)} \sum_{k=0}^{\infty} p_k\left(\frac{nx}{b_n}\right) f\left(\frac{k}{n} b_n\right)$$

with b_n a positive increasing sequence with the properties $\lim_{n\to\infty} b_n = \infty$ and $\lim_{n\to\infty} \frac{b_n}{n} = 0$. They also studied some approximation properties of the operators.

Now, let us define Choldowsky type generalization of the $q-$Favard-Szàsz operators as follows:

$$P_n^*(f;q;x) = \frac{E_q^{-\frac{[n]_q}{b_n}x}}{A(1)} \sum_{k=0}^{\infty} \frac{P_k\left(q; \frac{[n]_q}{b_n}x\right)}{[k]_q!} f\left(\frac{[k]_q}{[n]_q} b_n\right), \tag{2.1}$$

where $\{P_k(q;.)\} \geqslant 0$ is a $q-$Appell polynomial set which is generated by

$$A(u)\, e_q^{\frac{[n]_q}{b_n}xu} = \sum_{k=0}^{\infty} \frac{P_k\left(q; \frac{[n]_q}{b_n}x\right) u^k}{[k]_q!} \tag{2.2}$$

and $A(u)$ is defined by

$$A(u) = \sum_{k=0}^{\infty} a_k u^k.$$

3. weighted statistical approximation properties

In this section, we give Korovkin type weighted statistical approximation properties of the operators $P_n^*(f;q;x)$. Before proceeding further, let us give basic definition and notation on the concept of the statistical convergence which was introduced by Fast [10]. Let K be a subset of $\mathbb{N}$, the set of natural numbers. Then, $K_n = \{k \leqslant n : k \in K\}$. The natural density of K is defined by $\delta(K) = \lim_n \frac{1}{n}|K_n|$ provided that the limit exists, where $|K_n|$ denotes the cardinality of the set K_n. A sequence $x = (x_k)$ is called statistically convergent to the number $\ell \in \mathbb{R}$, denoted by $st - \lim x = \ell$. For each $\epsilon > 0$, the set $K_\varepsilon = \{k \in \mathbb{N} : |x_k - \ell| \geqslant \epsilon\}$ has a natural density zero, that is

$$\lim_{n\to\infty} \frac{1}{n} |\{k \leqslant n : |x_k - \ell| \geqslant \epsilon\}| = 0.$$

It is well know that every statistically convergence sequence is ordinary convergent, but the converse is not true. The concept of statistical convergence was firstly used in approximation theory by Gadjiev and Orhan [11]. They proved the Bohman−Korovkin type approximation theorem for statistical convergence.

A real function ρ_0 is called a weighted function if it is continuous on $\mathbb{R}$ and $\lim_{|x|\to 0} \rho_0(x) = \infty$, $\rho_0(x) \geqslant 1$ for all $x \in \mathbb{R}$, where $\rho_0(x) = 1 + x^2$. Let $B_{\rho_0}[0,\infty)$ be the set of all functions f defined on $[0,\infty)$ satisfying the condition $|f(x)| \leqslant M_f(1 + x^2)$, where M_f is a constant depending on f. By $C_{\rho_0}[0,\infty)$, we denote the subspace of all continuous function belonging to $B_{\rho_0}[0,\infty)$. Also, $C^*_{\rho_0}[0,\infty)$ be subspace of all continuous functions $f \in C_{\rho_0}[0,\infty)$, for which $\lim_{|x|\to\infty} \frac{f(x)}{1+x^2}$ is finite. The norm on $C^*_{\rho_0}[0,\infty)$ is defined as follows:

$$\parallel f \parallel_\rho = \sup_{x\in[0,\infty)} \frac{|f(x)|}{1+x^2}.$$

Now, we may begin the following lemma which is needed proving our main result.

Lemma 3.1. *For $n \in \mathbb{N}$, $x \in [0,\infty)$ and $0 < q < 1$, we have*

$$\begin{aligned}
&(3.1)\quad P_n^*(e_0;q;x) &=&\ 1,\\
&(3.2)\quad P_n^*(e_1;q;x) &=&\ x + \frac{D_q(A(1))E_q^{-\frac{[n]_q}{b_n}x} e_q^{\frac{q[n]_q}{b_n}x}}{A(1)} \frac{b_n}{[n]_q},\\
&(3.3)\quad P_n^*(e_2;q;x) &=&\ x^2 + \frac{E_q^{-\frac{[n]_q}{b_n}x} e_q^{q\frac{[n]_q}{b_n}x} \left[qD_q(A(q)) + D_q(A(1))\right]}{A(1)} \frac{xb_n}{[n]_q} + \frac{E_q^{-\frac{[n]_q}{b_n}x} e_q^{q\frac{[n]_q}{b_n}x} D_q^2(A(1))}{A(1)} \frac{b_n^2}{[n]_q^2},
\end{aligned}$$

where $e_i(x) = x^i$, $i = 0,1,2$.

Proof. By (2.2) and definition of q derivative, we obtain that

$$\sum_{k=0}^{\infty} \frac{p_k\left(q; \frac{[n]_q}{b_n}x\right)}{[k]_q!} = A(1)\, e_q^{\frac{[n]_q}{b_n}x}, \tag{3.4}$$

$$\sum_{k=0}^{\infty} \frac{p_k\left(q; \frac{[n]_q}{b_n}x\right)}{[k]_q!} [k]_q = A(1) \frac{[n]_q}{b_n} x e_q^{\frac{[n]_q}{b_n}x} + e_q^{q\frac{[n]_q}{b_n}x} D_q A(1) \tag{3.5}$$

$$\begin{aligned}
\sum_{k=0}^{\infty} \frac{p_k\left(q; \frac{[n]_q}{b_n}x\right)}{[k]_q!} [k]_q^2 &= D_q^2(A(1)) e_q^{\frac{q[n]_q}{b_n}x} + \frac{x[n]_q}{b_n} e_q^{\frac{q[n]_q}{b_n}x} \left[qD_q(A(q)) + D_q(A(1)\right] \\
&\quad + A(1) \frac{x^2[n]_q^2}{b_n^2} e_q^{\frac{[n]_q}{b_n}x}.
\end{aligned} \tag{3.6}$$

By using the relations (3.4)-(3.6), from (2.1), we obtain the results

$$P_n^*(e_0;q;x) = \frac{E_q^{-\frac{[n]_q}{b_n}x}}{A(1)} \sum_{k=0}^{\infty} \frac{p_k\left(q; \frac{[n]_q}{b_n}x\right)}{[k]_q!} = \frac{E_q^{-\frac{[n]_q}{b_n}x}}{A(1)} A(1)\, e_q^{\frac{[n]_q}{b_n}x} = 1,$$

$$\begin{aligned}
P_n^*(e_1;q;x) &= \frac{E_q^{-\frac{[n]_q}{b_n}x}}{A(1)} \sum_{k=0}^{\infty} \frac{p_k\left(q; \frac{[n]_q}{b_n}x\right)}{[k]_q!} \left(\frac{[k]_q}{[n]_q} b_n\right) \\
&= \frac{E_q^{-\frac{[n]_q}{b_n}x} b_n}{A(1)\,[n]_q} \sum_{k=0}^{\infty} \frac{p_k\left(q; [n]_q\, t\right)}{[k]_q!} [k]_q \\
&= \frac{E_q^{-\frac{[n]_q}{b_n}x} b_n}{A(1)\,[n]_q} \left(A(1) \frac{[n]_q}{b_n} x e_q^{\frac{[n]_q}{b_n}x} + e_q^{q\frac{[n]_q}{b_n}x} D_q A(1)\right) \\
&= x + \frac{D_q(A(1))E_q^{-\frac{[n]_q}{b_n}x} e_q^{\frac{q[n]_q}{b_n}x}}{A(1)} \frac{b_n}{[n]_q}
\end{aligned}$$

and

$$\begin{aligned}
P_n^*(e_2;q;x) &= \frac{E_q^{-\frac{[n]_q}{b_n}x}}{A(1)} \sum_{k=0}^{\infty} \frac{P_k\left(q; \frac{[n]_q}{b_n}x\right)}{[k]_q!} \frac{[k]_q^2}{[n]_q^2} b_n^2 \\
&= \frac{E_q^{-\frac{[n]_q}{b_n}x}}{A(1)} \frac{b_n^2}{[n]_q^2} \left\{ D_q^2(A(1)) e_q^{\frac{q[n]_q}{b_n}x} + e_q^{\frac{q[n]_q}{b_n}x} \frac{[n]_q}{b_n} x \left[qD_q(A(q)) + D_q(A(1)\right] + A(1) \frac{[n]_q^2}{b_n^2} x^2 e_q^{\frac{[n]_q}{b_n}x} \right\} \\
&= x^2 + \frac{E_q^{-\frac{[n]_q}{b_n}x} e_q^{q\frac{[n]_q}{b_n}x} \left[qD_q(A(q)) + D_q(A(1))\right]}{A(1)} \frac{xb_n}{[n]_q} + \frac{E_q^{-\frac{[n]_q}{b_n}x} e_q^{q\frac{[n]_q}{b_n}x} D_q^2(A(1))}{A(1)} \frac{b_n^2}{[n]_q^2}
\end{aligned}$$

Hence, the proof is completed. □

Theorem 3.2. *Assume that $q := (q_n)$, $0 < q_n < 1$ be a sequence satisfying the following conditions:*

$$st - \lim_n \frac{b_n}{[n]_q} = 0, \quad st - \lim_n q_n = b. \tag{3.7}$$

Let $f \in C[0,\infty) \cap E$, where $E = \{f \in C[0,\infty) : |f(x)| \leqslant \alpha e^{\beta x}$ for some $\alpha \in \mathbb{R}^+, \beta \in \mathbb{R}\}$. Then we have the following:

$$st - \lim_n \parallel P_n^*(f, q_n; .) - f \parallel_{\rho_0} = 0.$$

Proof. It is enough to prove that

$$st - \lim_n \parallel P_n^*(e_v, q_n; .) - e_v \parallel_{\rho_0} = 0$$

where $v = 0, 1, 2$.

From the equation (3.1), it is easy to obtain that

$$st - \lim_n \parallel P_n^*(e_0, q_n; .) - e_0 \parallel_{\rho_0} = 0.$$

By (3.2) and combining with $E_q^{-\frac{[n]_{q_n}}{b_n}} e_q^{\frac{q_n [n]_{q_n}}{b_n}} < 1$, we get

$$\parallel P_n^*(e_1, q_n; .) - e_1 \parallel_{\rho_0} \leqslant \frac{D_q(A(1))}{A(1)} \frac{b_n}{[n]_{q_n}} \parallel e_0 \parallel_{\rho_0} .$$

For $\epsilon > 0$, define the following sets:

$$\begin{aligned} \mathcal{M} &:= \left\{k : \parallel P_n^*(e_1, q_k; .) - e_1 \parallel_{\rho_0} \geqslant \epsilon\right\}, \\ \mathcal{M}_1 &:= \left\{k : \frac{D_q(A(1))}{A(1)} \frac{b_k}{[k]_{q_k}} \geqslant \epsilon\right\} \end{aligned}$$

such that $\mathcal{M} \subseteq \mathcal{M}_1$.

By (3.7), one can write the following

$$\delta \left\{k \leqslant n : \parallel P_n^*(e_1, q_k; .) - e_1 \parallel_{\rho_0} \geqslant \epsilon\right\} \leqslant \delta \left\{k \leqslant n : \frac{D_q(A(1))}{A(1)} \frac{b_k}{[k]_{q_k}} \geqslant \epsilon\right\}$$

Hence, we get

$$st - \lim_n \parallel P_n^*(e_1, q_n; .) - e_1 \parallel_{\rho_0} = 0. \tag{3.8}$$

By (3.3), one can see that

$$\begin{aligned} \parallel P_n^*(e_2, q_n; .) - e_2 \parallel_{\rho_0} \leqslant{} & \frac{q_n D_q(A(q)) + D_q(A(1))}{A(1)} \frac{b_n}{[n]_{q_n}} \parallel e_1 \parallel_{\rho_0} \\ & + \frac{D_q(A(1)) + D_q^2(A(1))}{A(1)} \frac{b_n^2}{[n]_{q_n}^2} \parallel e_0 \parallel_{\rho_0} \end{aligned}$$

Now, let $\epsilon > 0$ be given, we define the following sets:

$$\begin{aligned} \mathcal{V} &:= \left\{k : \parallel P_n^*(e_2, q_k; .) - e_2 \parallel_{\rho_0} \geqslant \epsilon\right\}, \\ \mathcal{V}_1 &:= \left\{k : \frac{q_k D_q(A(q)) + D_q(A(1))}{A(1)} \frac{b_k}{[k]_{q_k}} \geqslant \frac{\epsilon}{2}\right\}, \\ \mathcal{V}_2 &:= \left\{k : \frac{D_q(A(1)) + D_q^2(A(1))}{A(1)} \frac{b_k^2}{[k]_{q_k}^2} \geqslant \frac{\epsilon}{2}\right\}, \end{aligned}$$

such that $\mathcal{V} \subseteq \mathcal{V}_1 \cup \mathcal{V}_2$.

Thus, we obtain

$$\begin{aligned} \delta \left\{k \leqslant n : \parallel P_n^*(e_2, q_k; .) - e_2 \parallel_{\rho_0} \geqslant \epsilon\right\} \leqslant{} & \delta \left\{k \leqslant n : \frac{q_k D_q(A(q)) + D_q(A(1))}{A(1)} \frac{b_k}{[k]_{q_k}}\right\} \\ & + \delta \left\{k \leqslant n : \frac{D_q(A(1)) + D_q^2(A(1))}{A(1)} \frac{b_k^2}{[k]_{q_k}^2} \geqslant \frac{\epsilon}{2}\right\} \end{aligned} \tag{3.9}$$

Hence, (3.7) and (3.9) imply that

$$st - \lim_n \parallel P_n^*(e_2, q_n; .) - e_2 \parallel_{\rho_0} = 0. \tag{3.10}$$

□

4. Rates of statistical convergence

In this section, we give the rates of statistical convergence of the operators $P_n^*(f;q;x)$ by means of modulus of continuity with the help of functions from Lipschitz class.

$C_B[0,\infty)$ denotes of all real valued continuous bounded functions. The modulus of continuity for the function $f\in C_B[0,\infty)$ is defined as

$$\omega(f,\delta)_{\rho_0}=\sup_{\substack{|x-y|\leqslant\delta\\ x,y\in[0,\infty)}}\frac{|f(x)-f(y)|}{1+x^{2+\lambda}}$$

where $\omega(f,\delta)_{\rho_0}$ for $\delta>0$, $\lambda\geqslant 0$ satisfy the following conditions: for every $f\in C_B[0,\infty)$.

$$\lim_{\delta\to\infty}\omega(f,\delta)_{\rho_0}=0$$

and

$$|f(x)-f(y)|\leqslant\omega(f,\delta)_{\rho_0}\left(\frac{|x-y|}{\delta}+1\right). \tag{4.1}$$

Now, we prove the following theorem for the rate of pointwise convergence of the operators $P_n^*(f;q;x)$ to the function $f(x)$ by means of modulus of continuity.

Theorem 4.1. *$q:=(q_n)$ be sequence and $x\in[0,\infty)$, then we have*

$$|P_n^*(f;q;x)-f(x)|\leqslant 2\omega(f,\sqrt{\delta_n})_{\rho_0},$$

for all $f\in C_B[0,\infty)\cap E$, where

$$\delta_n=\frac{q_nD_q(A(q))+D_q(A(1))}{A(1)}\frac{b_n}{[n]_{q_n}}\parallel e_1\parallel_{\rho_0}+\frac{D_q^2(A(1))}{A(1)}\frac{b_n^2}{[n]_{q_n}^2}\parallel e_0\parallel_{\rho_0} \tag{4.2}$$

Proof. By the linearity and positivity of the operators $P_n^*(f;q;x)$, one can write

$$\begin{aligned}
|P_n^*(f;q;x)-f(x)| &\leqslant \frac{E_q^{-\frac{[n]_q}{b_n}x}}{A(1)}\sum_{k=0}^{\infty}\frac{p_k\left(q;\frac{[n]_q}{b_n}x\right)}{[k]_q!}\left|f\left(\frac{[k]_q}{[n]_q}b_n\right)-f(x)\right|\\
&\leqslant \frac{E_q^{-\frac{[n]_q}{b_n}x}}{A(1)}\sum_{k=0}^{\infty}\frac{p_k\left(q;\frac{[n]_q}{b_n}x\right)}{[k]_q!}\omega(f,\delta)\left\{\frac{1}{\delta}\left|\frac{[k]_q}{[n]_q}b_n-x\right|+1\right\}\\
&= \left\{\frac{1}{\delta}\frac{E_q^{-\frac{[n]_q}{b_n}x}}{A(1)}\sum_{k=0}^{\infty}\frac{p_k\left(q;\frac{[n]_q}{b_n}x\right)}{[k]_q!}\left|\frac{[k]_q}{[n]_q}b_n-x\right|+1\right\}\omega(f,\delta).
\end{aligned}$$

If we apply Cauchy-Schwarz inequality for sums, we obtain

$$\sum_{k=0}^{\infty}\frac{p_k\left(q;\frac{[n]_q}{b_n}x\right)}{[k]_q!}\left|\frac{[k]_q}{[n]_q}b_n-x\right|^2\leqslant\left(\sum_{k=0}^{\infty}\frac{p_k\left(q;[n]_q\,t\right)}{[k]_q!}\right)^{1/2}\left(\sum_{k=0}^{\infty}\frac{p_k\left(q;\frac{[n]_q}{b_n}x\right)}{[k]_q!}\left(\frac{[k]_q}{[n]_q}b_n-x\right)^2\right)^{1/2}.$$

Using above inequality and by (3.1), we have

$$\begin{aligned}
|P_n^*(f;q;x)-f(x)| &\leqslant \omega(f,\delta)\left\{1+\frac{1}{\delta}\left[\frac{E_q^{-\frac{[n]_q}{b_n}x}}{A(1)}\sum_{k=0}^{\infty}\frac{p_k\left(q;\frac{[n]_q}{b_n}x\right)}{[k]_q!}\left(\frac{[k]_q}{[n]_q}b_n-x\right)^2\right]^{1/2}\right\}\\
&= \omega(f,\delta)\left\{1+\frac{1}{\delta}\left[\frac{E_q^{-\frac{[n]_q}{b_n}x}}{A(1)}\sum_{k=0}^{\infty}\frac{p_k\left(q;\frac{[n]_q}{b_n}x\right)}{[k]_q!}\left(\frac{[k]_q}{[n]_q}b_n-x\right)^2\right]^{1/2}\right\}\\
&= \omega(f,\delta)\left\{1+\frac{1}{\delta}\left[P_n^*\left(s-e_1\right)^2;q;x)\right]^{1/2}\right\}.
\end{aligned}$$

From Lemma 3.1 and the fact $E_q^{-\frac{[n]_{q_n}}{b_n}}e_q^{\frac{q_n[n]_{q_n}}{b_n}}<1$, we have

$$P_n^*\left((s-e_1)^2;q_n;x\right)\leqslant\frac{q_nD_q(A(q))+D_q(A(1))}{A(1)}\frac{b_n}{[n]_{q_n}}\parallel e_1\parallel_{\rho_0}+\frac{D_q^2(A(1))}{A(1)}\frac{b_n^2}{[n]_{q_n}^2}\parallel e_0\parallel_{\rho_0}. \tag{4.3}$$

By (4.3) and we choose $\delta=\sqrt{\delta_n}$, we get

$$|P_n^*(f;q;x)-f(x)|\leqslant 2\omega(f,\sqrt{\delta_n})_{\rho_0}.$$

This step concludes the proof. ☐

We know that a function $f \in C[0, \infty)$ is in $Lip_M(\alpha)$ on F, $0 < \alpha \leqslant 1$, F is a any bounded subset of the interval $[0, \infty)$ if it is satisfies the conditions

$$|f(y) - f(x)| \leqslant M|x - y|^{\alpha},\ y \in [0, \infty) \text{ and } x \in F, \tag{4.4}$$

where M is a constant depending only α and f.

Theorem 4.2. *Let $f \in C[0, \infty) \cap Lip_M(\alpha)$, $0 < \alpha \leqslant 1$. Then, we get*

$$|P_n^*(f; q_n; x) - f(x)| \leqslant M \left(\delta_n^{\alpha/2} + d^{\alpha}(x, F) \right),\ x \in [0, \infty),$$

where M is a constant depending only α and f and $d(x, F)$ is the distance between x and F defined by $d(x, F) = \inf\{|x - t| : t \in F\}$.

Proof. Let $\overline{F}$ be the closure of F in $[0, \infty)$. The there exists al leat point $x_0 \in \overline{F}$ such that $d(x, F) = |x - x_0|$. By our assumption and monotonicity of $P_n^*(f; q_n; x)$, one can write the following

$$\begin{aligned} |P_n^*(f; q_n; x) - f(x)| &\leqslant P_n^*(|f(t) - f(x)|; q_n; x) + P_n^*(|f(x) - f(x_0)|; q_n; x) \\ &\leqslant M\{P_n^*(|t - x|^{\alpha}; q_n; x) + |x - x_0|^{\alpha}\}. \end{aligned}$$

Next, we applying the Hölder inequality with $p = \frac{1}{\alpha}, q = \frac{\alpha}{2-\alpha}$ and we get

$$|P_n^*(f; q_n; x) - f(x)| \leqslant M\{P_n^*\left[|t - x|^2; q_n; x\right]^{\alpha/2} + |x - x_0|^{\alpha}\}.$$

This step conclude the proof. ☐

Now, we obtain local direct estimate of the operators $P_n^*(f; q; x)$ using the Lipschitz-type maximal function of order α introduce Lenz [17] as

$$\omega_{\alpha}(f, x) = \sup_{t \neq x,\ t \in [0, \infty)} \frac{|f(t) - f(x)|}{|t - x|^{\alpha}},\ x \in [0, \infty) \text{ and } \alpha \in (0, 1]. \tag{4.5}$$

Theorem 4.3. *Let $f \in Lip_M(\alpha)$, $0 < \alpha \leqslant 1$, then we have*

$$|P_n^*(f; q_n; .) - f(.)| \leqslant \omega_{\alpha}(f, x)\delta_n^{\alpha/2},$$

where δ_n defined in (5.1).

Proof. From (4.5), we obtain

$$\begin{aligned} |P_n^*(f; q_n; x) - f(x)| &\leqslant P_n^*(|f(t) - f(x)|, q; x) \\ &\leqslant \omega_{\alpha}(f, x) P_n^*(|t - x|^{\alpha}, q; x). \end{aligned}$$

Again for $p = \frac{1}{\alpha}, q = \frac{\alpha}{2-\alpha}$, applying the Hölder inequality, we get

$$\begin{aligned} |P_n^*(f; q_n; x) - f(x)| &\leqslant P_n^*(|f(t) - f(x)|, q; x) \\ &\leqslant \omega_{\alpha}(f, x)\{P_n^*(e_1 - x)^2, q; x)\}^{\alpha/2} = \omega_{\alpha}(f, x)\delta_n^{\alpha/2}. \end{aligned}$$

Hence, we get the desired result. ☐

5. Local Approximation

In this section, we state the local approximation theorem of the operators $P_n^*(f; q; x)$. Let $C_B[0, \infty)$ be the space of all real valued continuous bounded functions f on $[0, \infty)$ with the norm $\| f \| = \sup\{|f(x)| : x \in [0, \infty)\}$. The K-functional of f is defined by

$$K_2(f; \delta) = \inf_{g \in W^2} \{\|f - g\| + \delta\|g''\|\},$$

where $\delta > 0$ and $W^2 = \{g \in C_B[0, \infty) : g', g'' \in C_B[0, \infty)\}$. By Devore-Lorentz [5, p. 177], there exists an absolute constant $C > 0$ such that

$$K_2(f, \delta) \leqslant C\omega_2\left(f, \sqrt{\delta}\right) \tag{5.1}$$

where

$$\omega_2\left(f, \sqrt{\delta}\right) = \sup_{0 < h \leqslant 0} \sup_{x \in [0, \infty)} |f(x + 2h) - 2f(x + h) + f(x)|$$

is the second order modulus of smoothness of f. Moreover,

$$\omega(f,\delta)=\sup_{0<h\leqslant 0}\ \sup_{x\in[0,\infty)}|f(x+h)-f(x)|$$

denotes the modulus of continuity of f.

Now, we give the direct local approximation theorem for the operators $P_n^*(f;q;x)$.

Theorem 5.1. *Let $q\in(0,1)$. We have*

$$|P_n^*(f;q;x)-f(x)|\leqslant K\omega_2(f,\varphi_n)+\omega\left(f,\frac{D_q(A(1))}{A(1)}\frac{b_n}{[n]_q}\right)$$

$\forall x\in[0,\infty)$, $f\in C_B[0,\infty)$, where K is a positive constant and

$$\varphi_n=\frac{q_nD_q(A(q))+D_q(A(1))}{A(1)}\frac{b_n}{[n]_{q_n}}+\frac{A(1)D_q^2(A(1))+[D_q(A(1))]^2}{A^2(1)}\frac{b_n^2}{[n]_{q_n}^2}$$

Proof. Let us define the following operators

$$\widetilde{P}_n^*(f;q;x)=P_n^*(f;q;x)-f\left(x+\frac{D_q(A(1))E_q^{-\frac{[n]_q}{b_n}x}e_q^{\frac{q[n]_q}{b_n}x}}{A(1)}\frac{b_n}{[n]_q}\right)+f(x), \tag{5.2}$$

$x\in[0,\infty)$. The operators $\widetilde{P}_n^*(f;q;x)$ are linear. Thus, we have the following:

$$\widetilde{P}_n^*(s-x;q;x)=0, \tag{5.3}$$

(see Lemma 3.1). Let $g\in W^2$, from Taylor's expansion

$$g(s)=g(x)+g'(x)(s-x)+\int_x^s(s-u)g''(x)du,$$

$s\in[0,\infty)$ and (5.3) we obtain

$$\widetilde{P}_n^*(g;q;x)=g(x)+\widetilde{P}_n^*\left(\int_x^s(s-u)g''(x)du\right).$$

By (5.2), we have the following

$$\begin{aligned}
|\widetilde{P}_n^*(g;q;x)-g(x)| &\leqslant \left|P_n^*\left(\int_x^s(s-u)g''(u)du\right)\right|\\
&\quad+\left|\int_x^{x+\frac{D_q(A(1))E_q^{-\frac{[n]_q}{b_n}x}e_q^{\frac{q[n]_q}{b_n}x}}{A(1)}\frac{b_n}{[n]_q}}\left(x+\frac{D_q(A(1))E_q^{-\frac{[n]_q}{b_n}x}e_q^{\frac{q[n]_q}{b_n}x}}{A(1)}\frac{b_n}{[n]_q}-u\right)g''(u)du\right|\\
&\leqslant P_n^*\left(\left|\int_x^s(s-u)g''(u)du\right|,x\right)\\
&\quad+\int_x^{x+\frac{D_q(A(1))E_q^{-\frac{[n]_q}{b_n}x}e_q^{\frac{q[n]_q}{b_n}x}}{A(1)}\frac{b_n}{[n]_q}}\left|x+\frac{D_q(A(1))E_q^{-\frac{[n]_q}{b_n}x}e_q^{\frac{q[n]_q}{b_n}x}}{A(1)}\frac{b_n}{[n]_q}-u\right||g''(u)|\,du\\
&\leqslant \left[P_n^*(s-x)^2+\left(\frac{D_q(A(1))E_q^{-\frac{[n]_q}{b_n}x}e_q^{\frac{q[n]_q}{b_n}x}}{A(1)}\frac{b_n}{[n]_q}\right)^2\right]\|g''\| \qquad (5.4)
\end{aligned}$$

Therefore, from (5.4), we obtain

$$|\widetilde{P}_n^*(g;q;x)-g(x)|\leqslant\varphi_n. \tag{5.5}$$

By (2.1), (3.1) and (5.2), we get

$$\begin{aligned}
|\widetilde{P}_n^*(f;q;x)| &\leqslant P_n^*(f;q;x)+2\|f\|\\
&\leqslant \|f\|P_n^*(1;q;x)+2\|f\|\\
&\leqslant 3\|f\| \qquad (5.6)
\end{aligned}$$

and by (5.2), (5.5) and (5.6)

$$\begin{aligned} \left|P_n^*(f;q;x)-f(x)\right| &\leqslant \left|\widetilde{P}_n^*(f-g;q;x)-(f-g)(x)\right|+\left|\widetilde{P}_n^*(g;q;x)-g(x)\right| \\ &\quad+\left|f\left(x+\frac{D_q(A(1))E_q^{-\frac{[n]_q}{b_n}x}e_q^{\frac{q[n]_q}{b_n}x}}{A(1)}\frac{b_n}{[n]_q}\right)-f(x)\right| \\ &\leqslant 4\|f-g\|+\varphi_n\|g''\|. \end{aligned}$$

From (5.1), one can see that

$$\left|P_n^*(f;q;x)-f(x)\right|\leqslant K\omega_2(f,\varphi_n)+\omega\left(f,\frac{D_q(A(1))}{A(1)}\frac{b_n}{[n]_q}\right).$$

and this concludes the proof. ☐

References

[1] A. Aral, V. Gupta, On the Durrmeyer type modification of the $q-$Baskakov type operators, Nonlinear Anal. 72(3) (2010) 1171–1180.

[2] Ç. Atakut, İ. Büyükyazıcı , The rate of convergence of the $q-$analogue of Favard Szàsz type operators, J. Comput. Anal. Appl. 13 (4) (2011) 673–682.

[3] Z. Finta, V. Gupta, Approximation by $q-$Durrmeyer operators, J. Appl. Math. Comput. 29 (1–2) (2009) 401–415.

[4] İ. Büyükyazıcı, Ç. Atakut, On Stancu type generalization of $q-$Baskakov operators, Math. Comput. Modelling 52(5) (2010) 752–759.

[5] R.A. DeVore, G.G. Lorentz, Constr. Approx. Springer, Berlin, 1993.

[6] G. M. Phillips, Bernstein polynomials based on the $q-$integers, Ann. Numer. Math. 4 (1997), no. 1-4, 511-518.

[7] Ç. Atakut, H. Tanberkan, S. K. Serenbay and İ. Büyükyazıcı, Approximation by Chlodowsky type Jakimovski-Leviatan operators, J. Comput. Appl. Math., 259, 153–163 (2014).

[8] A. Aral, V. Gupta and R. P. Agarwal, Applications of $q-$calculus in operator theory, Berlin, Springer, 2013.

[9] A. Jakimovski, D. Leviatan, Generalized Szàsz operators for the approximation in the infinite interval, Mathematica (Cluj) 34 (1969) 97–103.

[10] H. Fast, Sur la convergence statistique, Colloq. Math. Studia Math. 2 (1951) 241-244.

[11] A.D. Gadjiev, C. Orhan, Some approximation properties via statistical convergence, Rocky Mountain J. Math. 32 (2002) 129-138.

[12] Ö. Dalmanoğlu, O. Doğru, On statistical approximation properties of Kantorovich type $q-$Bernstein operators, Math. Comput. Modelling 52(2010) 760–771.

[13] C. Radu, Statistical approximation properties of Kantorovich operators based on q-integers, Creative Math. and Inf, 17 (2008) 75–84.

[14] V. Gupta, C. Radu, Statistical approximation properties of $q-$Baskakov-Kantorovich operators, Cent. Eur. J. Math. 7 (2009) (4) 809-818

[15] M. Örükçü, O. Doğru, Weighted statistical approximation by Kantorovich type $q-$Szàsz-Mirakjan operators, Appl. Math. Comput. 217 (2011) 20 7913-7919.

[16] M. Mursaleen, A. Khan, Statistical approximation properties of modified $q-$Stancu-Beta operators, Bull. Malays. Math. Sci. Soc.(2) 36.3 (2013) 683–690.

[17] B. Lenze, Bernstein-Baskakov-Kantorovic operators and Lipschitz-type maximal functions, in Approximation Theory (Kecskemet, 1990) 469496, Colloq. Math. Soc. Janos Bolyai, 58 North-Holland, Amsterdam.

[18] M. Mursaleen, V. Karakaya, M. Ertürk, F. Gürsoy, Weighted statistical convergence and its application to Korovkin type approximation theorem, Appl. Math. Comput (218) 18 (2012): 9132–9137.

[19] M. Mursaleen, A. Alotaibi, Statistical summability and approximation by de la Valle-Poussin mean, Appl. Math. Lett. (24) 3 (2011) 320–324.

[20] M. Mursaleen, A. Alotaibi, Statistical lacunary summability and a Korovkin type approximation theorem, Ann. Univ. Ferrara. 57(2) (2011) 373–381.

[21] V.Karakaya, A.Karaisa, Korovkin type approximation theorems for weighted $\alpha\beta$-statistical convergence, Bull. Math. Sci. DOI 10.1007/s13373-015-0065-y,(2015).